

If you have any questions or concerns, please consult with a pharmacist or a physician as needed.

Check (1) Do **all** of the following apply to you?

- ☐ Are you unable to experience an erection of sufficient hardness or of sustained duration to enjoy sexual intercourse?
- ☐ Are you an adult male (18 years or older)?

NO

If you answered "No"
to at least one question,
**you cannot use
this medication.**

Yes

Check (2) Do **any** of the following apply to you?

- ☐ Have you ever experienced an allergic reaction from this medication or from the ingredients of this medication?
- ☐ Do you use nitrates or similar substances commonly used for treating heart disease such as angina, such as nitroglycerin (Millisrol, etc.), isosorbide dinitrate (Nitorol, Frandol, etc.), and nicorandil (SIGMART, etc.)?
- ☐ Do you use amyl nitrites (including recreational poppers, etc.), etc.?
- ☐ Do you use riociguat (Adempas, etc.), ketoconazole (for oral administration, not available in Japan), itraconazole (ITRIZOLE, etc.), or clarithromycin (Clarice, Klaricid, etc.)?

Yes

If you answered "Yes"
to at least one question,
**you cannot use
this medication.**

You are unable to receive treatment
using this medication, so please consult
with a physician about a different form
of treatment.

NO

Check (3) Do **any** of the following apply to you?

- ☐ Are you suspected of having sexual dysfunction due to factors such as cardiovascular system-related dysfunction?
- ☐ Have you ever experienced unstable angina or angina during sexual intercourse?
- ☐ Do you experience an irregular pulse despite taking the appropriate medication?
- ☐ Do you have low blood pressure (blood pressure < 90/50 mmHg)?
- ☐ Do you experience high blood pressure despite taking medication to lower your blood pressure (blood pressure after resting > 170/100 mmHg)?
- ☐ Have you experienced a myocardial infarction within the past three months?
- ☐ Have you experienced a cerebral infarction or cerebral hemorrhage within the past six months?
- ☐ Has a physician indicated that you may have a liver abnormality?
- ☐ Has a physician indicated that you may have a kidney abnormality?
- ☐ Have you ever been diagnosed with the following?
• Retinal pigmentary degeneration

Yes

If you answered "Yes"
to at least one question,
**you cannot use
this medication.**

Sexual intercourse involves risk of heart-related issues. If you have suspected heart or circulatory system-related dysfunction, please make sure to consult with a physician beforehand. You may be at risk for myocardial infarction or a similar condition. Also, you cannot use this medication if you have severe heart dysfunction, moderate to severe kidney dysfunction, or you have been diagnosed with retinal pigmentary degeneration. Please consult with a physician about a different form of treatment.

NO

Check (4) Do **any** of the following apply to you?

- ☐ Are you receiving any treatment or taking medication prescribed by a physician?
- ☐ Have you ever experienced an allergic reaction from medication, etc.?
- ☐ Do you use any of the following medications?
• Benign prostatic hyperplasia therapeutics or drugs that lower blood pressure (alpha blockers (doxazosin etc.) amlodipine, etc.)
• Anti-HIV agents (ritonavir, saquinavir, etc.)
• Antibacterial agents (rifampicin, etc.)
• Antiepileptic drugs (phenytoin, phenobarbital, etc.)
- ☐ Have you ever been diagnosed with the following?
• Priapism (sustained erection lasting at least 4 hours)
• Blood-related disease (sickle cell anemia, multiple myeloma, leukemia, etc.)
• Hemorrhagic disease, peptic ulcer
• Genital deformation (bending, genital fibrosis, Peyronie's disease, etc.)
- ☐ Are you 65 years or older?
• Have you never received a medical examination to test kidney function?
• Has a recent medical examination indicated that you have decreased kidney function (Ccr (Creatine Clearance) of 50 mL/min or less)?
- ☐ Are you receiving any other treatment for erectile dysfunction?
- ☐ Do you experience intense difficulty breathing, chest pains, or similar symptoms following light or moderate exercise (for example, when walking quickly for around 20 minutes, when quickly ascending one flight of stairs, etc.)?

Yes

If you answered "Yes"
to at least one question,
it is highly likely that
**you cannot use
this medication,
so please consult
with a physician or
pharmacist.**

Please consult with a physician about
taking this medication during a medical
examination at a medical facility.

If a physician advises
that you are able to use
this medication

NO

Before using this medication, please check with a pharmacist that you will adhere to the following items.

After confirming the following points ☒ please check the box.

- ☐ Please refrain from taking grapefruit juice, etc., which affect this drug.
- ☐ If you do not experience any effects after 4 to 8 uses, please consult with a physician or pharmacist. (Erectile dysfunction may be associated with conditions such as heart disease, high blood pressure, and diabetes, so please immediately consult with a physician or pharmacist.)
- ☐ If you experience symptoms such as angina following use of this medication, please make sure to consult with a physician or pharmacist regarding your use of this medication during a medical examination at a medical facility.
- ☐ After taking this medication, do not work at heights or operate vehicles or heavy machinery. (Use of this medication can result in symptoms such as dizziness or blurred vision.)

I have confirmed the above.

After receiving an instruction from a pharmacist, please make sure that you have carefully read the Informational Materials for Users before purchasing the medication. Also, even following purchase, make sure that you thoroughly understand the contents of the Package Insert before use. Please consult with a pharmacist if you require any help.

