

Please keep this leaflet and drug together at all times and read carefully before using.

EVE A EX

Pain Reliever/Fever Reducer

Designated Class 2 OTC drug

⚠ CAUTIONS FOR USE**⊗ Contraindications**

(Failure to observe the following may cause the present symptoms to worsen or produce adverse drug reactions and/or accidents.)

- The following people must not take this drug:**
 - Those with a previous allergic reaction to this drug or ingredients of the drug
 - Those who have ever suffered from asthma after taking this drug or other drugs for the relief of pain and fever or after taking other cold medicines
 - Children under 15 years of age
 - Pregnant women whose due date is within 12 weeks
- Do not take this drug in combination with any of the following drugs:**
Other pain reliever and fever reducers, cold medicines, sedatives or motion sickness drugs
- Do not drive or operate machinery after taking this drug.**
(Sleepiness may occur.)
- Do not drink alcohol before or after taking this drug.**
- Do not use this drug for a prolonged period of time.**

**When to consult a doctor**

- The following people should consult a doctor, dentist, pharmacist or registered salesperson before taking this drug:**
 - Those who are under the treatment of a doctor or dentist
 - Pregnant women or those who may become pregnant
 - Breastfeeding women
 - Elderly people
 - Those with any previous allergic reactions to drugs or other things
 - Those who were diagnosed with any of the following diseases:
Heart disease, kidney disease, liver disease, systemic lupus erythematosus or mixed connective-tissue disease
 - Those who have suffered from any of the following diseases:
Gastric and duodenal ulcers, ulcerative colitis or Crohn's disease

- The following symptoms may be adverse drug reactions. If any of these symptoms occur, STOP TAKING the drug immediately and consult a doctor, pharmacist or registered salesperson. Show them this leaflet.**

Related areas	Symptoms
Skin	Rash, redness, itching or purpura
Digestive system	Nausea, vomiting, loss of appetite, stomach discomfort, stomachache, oral ulcer, heartburn, heavy stomach feeling, gastrointestinal bleeding, abdominal pains, diarrhea or bloody stool
Central nervous system	Dizziness/vertigo
Circulatory system	Palpitations
Respiratory system	Breath shortness
Other	Blurred vision, ringing in the ears, swelling, nose bleeding, gum bleeding, uncontrolled bleeding, bleeding, back pain, excessive fall in temperature or malaise

In rare cases the following serious symptoms could occur. In such an event, immediately consult a doctor.

Symptoms' name	Symptoms
Shock (Anaphylaxis)	Itching of skin, hives, hoarseness, sneezing, itching of throat, tightness in the chest, palpitations, consciousness clouding, etc. which develop soon after taking the drug
Mucocutaneous ocular syndrome (Stevens-Johnson syndrome), Toxic Epidermal necrolysis	Persistent or acute worsening of high fever, bloodshot eyes, eye discharge, lip erosion, sore throat, widespread rash, redness, etc.
Liver dysfunction	Fever, itching, rash, jaundice (skin and whites of the eyes turn yellow), brown urine, general malaise, loss of appetite, etc.
Kidney disorder	Fever, rash, reduction of urine output, generalized swelling, general malaise, joint pain, diarrhea, etc.

(Continued on the next page)

Symptoms' name	Symptoms
Aseptic meningitis	Severe headache with neck muscle rigidity, fever, nausea, vomiting, etc. (These symptoms are mainly reported in patients undergoing treatment for systemic lupus erythematosus or mixed connective-tissue disease.)
Asthma	Wheezing or crowing sound when breathing, breathlessness, etc.
Aplastic anemia	Purpura, nose bleeding, gum bleeding, fever, pale skin, pale mucosa, tiredness, palpitations, breathlessness, giddy feeling accompanied by queasy feeling, bloody urine, etc.
Agranulocytosis	Sudden high fever, chill, sore throat, etc.

3. You may develop the following symptoms after taking this drug. If the symptoms continue or worsen, stop taking the drug and consult a doctor, pharmacist or registered salesperson. Show them this leaflet.

Constipation or sleepiness

4. If the symptoms do not improve after 3 or 4 doses, stop taking this drug and consult a doctor, dentist, pharmacist or registered salesperson. Show them this leaflet.

■ Indications

- Relief of pain associated with menstruation, headache, toothache, sore throat, joints, muscles, neuralgia, lower-back, stiff-shoulders, tooth extraction, bruising, earache, bone fracture, sprains and wounds
- Fever reduction, relief from chills due to fever

■ Dosage and Directions

Take the following dose up to 2 times a day, at intervals of at least 6 hours, avoid taking on an empty stomach. This drug should be taken with either cold or lukewarm water.

- **Adults (15 years and over):** 2 tablets
- **Children under 15 years of age:** DO NOT TAKE

<Cautions>

- (1) Strictly observe the dosage and directions.
- (2) Removing tablets from blister pack:
As shown in the figure on the right, push firmly on the blister until the tablet pierces the seal on the reverse side, then remove the tablet (Taking the tablet without removing it from the blister pack may cause esophageal mucosal injury).



■ Ingredients

Two tablets contain:

- **Ibuprofen** - 200 mg
- **Allylisopropylacetylurea** - 60 mg
- **Anhydrous Caffeine** - 80 mg

Additives: Anhydrous silicic acid, cellulose, croscarmellose sodium, polyvinyl alcohol·acrylic acid·methyl methacrylate copolymer, talc, magnesium stearate, hydroxypropyl cellulose, hypromellose, macrogol and titanium oxide

■ Storage and Handling Cautions

- (1) Store this drug in a cool, dry place away from sunlight and moisture.
- (2) Keep out of reach of children.
- (3) Do not transfer this drug to a different container as it may cause misuse or deterioration.
- (4) Do not use this drug after the expiration date.

製造販売元  **エスエス製薬株式会社**
〒163-1488 東京都新宿区西新宿3-20-2
<https://www.ssp.co.jp/>

The marketing of this drug has been approved for domestic sales in accordance with the PMD Act (The Pharmaceuticals, Medical Devices and other therapeutic products Act) in Japan.