

Please read the instructions carefully before using.

新エスタック®顆粒

かぜの諸症状（のどの痛み・熱）の緩和に **第②類医薬品**

※ この医薬品は「使用上の注意」をよく読んでお使いください。

アレルギー体質の方は、必ず薬剤師、登録販売者にご相談ください。



NEW S. TAC GRANULES

Designated Class 2 OTC drug

Cold medicine

It contains Kakkonto-ka-kikyo extract

Granules easy to swallow and easy to solve

- NEW S. TAC GRANULES is a cold medicine which combines Western medicinal ingredients effective for cold symptoms and “Kakkonto-ka-kikyo extract” that is composed with traditional cold medicine Kakkonto added with Kikyo.
- NEW S. TAC GRANULES is a formula of granules easy to swallow and easy to solve.

CAUTIONS FOR USE

CONTRAINDICATIONS

(Failure to observe the following may cause the present symptoms to worsen or produce adverse drug reactions and/or accidents.)

1. The following people must not take this drug:

- (1) Those with a previous allergic reaction to this drug or ingredients of the drug
- (2) Those who have ever suffered from asthma after taking this drug or other drugs for cold remedies or for relief of pain and fever
- (3) Children under 12 years of age

2. Do not take this drug in combination with any of the following drugs:

Other cold medicines, pain relievers/fever reducers, sedatives, cough and expectorant medicines, antihistamine-containing oral drugs (e.g. oral nasal inflammation drugs, motion sickness drugs, anti-allergy drugs, hypnotics and sedatives, etc.)

3. Do not drive or operate machinery after taking this drug.

(Sleepiness, etc. may occur.)

4. Breastfeeding women should not take this drug. If unavoidable, refrain from breastfeeding while taking this drug.

5. Do not drink alcohol before or after taking this drug.

6. Do not use this drug for a prolonged period of time.

WHEN to consult a doctor

1. The following people should consult a doctor, pharmacist or registered salesperson before taking this drug:

- (1) Those who are under the treatment of a doctor or dentist

®: Trademark



SSP Co., Ltd.

- (2) Pregnant women or those who may become pregnant
 (3) Elderly people
 (4) Those with any previous allergic reactions to drugs or other things
 (5) Those with any of the following symptoms: High fever or urination difficulties
 (6) Those who were diagnosed with any of the following diseases:
 Disrupted thyroid function, diabetes, heart disease, hypertension, liver disease, kidney disease, gastric and duodenal ulcers, glaucoma, respiratory functional disorder, obstructive sleep apnoea syndrome (OSAS), obesity or phenylketonuria

2. The following symptoms may be adverse drug reactions. If any of these symptoms occur, STOP TAKING the drug immediately and consult a doctor, pharmacist or registered salesperson. Show them this leaflet.

Related areas	Symptoms
Skin	Rash, redness, itching
Digestive system	Nausea, vomiting, loss of appetite
Central nervous system	Dizziness/vertigo
Urinary system	Urination difficulties
Other	Excessive fall in temperature

In rare cases, the following serious symptoms could occur. In such an event, immediately consult a doctor.

Symptom's name	Symptoms
Shock (Anaphylaxis)	Itching of skin, hives, hoarseness, sneezing, itching of throat, tightness in the chest, palpitations, consciousness clouding, etc. which develop soon after taking the drug
Mucocutaneous ocular syndrome (Stevens-Johnson syndrome), Toxic Epidermal necrolysis, Acute generalized exanthematous pustulosis (AGEP)	Persistent or acute worsening of high fever, bloodshot eyes, eye discharge, lip erosion, sore throat, widespread rash, redness, redness of skin with small pimples, general malaise, loss of appetite, etc.
Drug-induced hypersensitivity syndrome	Redness of the skin over a wide range, generalized rash, fever, dullness, swelling of lymph nodes (neck, underarm, base of hip, etc.), etc.
Liver dysfunction	Fever, itching, rash, jaundice (skin and whites of the eyes turn yellow), brown urine, general malaise, loss of appetite, etc.
Kidney disorder	Fever, rash, reduction of urine output, generalized swelling, general malaise, joint pain (painful joints), diarrhea, etc.
Interstitial pneumonia	Shortness of breath, difficulty breathing when going up stairs or performing strenuous activity, dry cough, fever or other symptoms, which happen suddenly or continuously

(Continued on the next page)

(Continued from the previous page)

Symptom's name	Symptoms
Asthma	Wheezing or crowing sound when breathing breathlessness, etc.
Aplastic anemia	Purpura, nose bleeding, gum bleeding, fever, pale skin, pale mucosa, tiredness, palpitations, breathlessness, giddy feeling accompanied by queasy feeling, bloody urine, etc.
Agranulocytosis	Sudden high fever, chill, sore throat, etc.
Respiratory depression	Shortness of breath or difficulty breathing, etc.

- 3. You may develop the following symptoms after taking this drug. If the symptoms continue or worsen, stop taking the drug and consult a doctor, pharmacist or registered salesperson. Show them this leaflet.**

Constipation, dry mouth or sleepiness

- 4. If the symptoms do not improve after 5 or 6 doses, stop taking this drug and consult a doctor, pharmacist or registered salesperson. Show them this leaflet.**

Indications

Alleviation of various common cold symptoms (fever, chills, headache, runny nose, nasal congestion, sneezing, cough, sore throat, joint pain and muscle pain)

Dosage and Directions

Take the following dose 3 times a day, preferably within 30 minutes of a meal. This drug should be taken with either cold or lukewarm water.

Age	Adult (15 years or over)	12-14 years	Under 12 years
One dose	One sachet	Two-thirds of a sachet	Do not take

<Cautions regarding dosage and directions>

- (1) Strictly observe the dosage and directions.
- (2) Children must be supervised by a parent or guardian when taking this drug.

Ingredients

One sachet (2.0 g) contains:

- Kakkonto-ka-kikyo extract - 1,000 mg
(Extracted from the infused liquid of the following:
pueraria root 0.53 g, ephedra herb 0.27 g, cinnamon bark 0.2 g, peony root 0.2 g, jujube 0.27 g, ginger 0.07 g, glycyrrhiza 0.13 g and platycodon root 0.27 g)
- Acetaminophen - 240 mg
- Dihydrocodeine Phosphate - 5 mg
- Chlorpheniramine Maleate - 2.5 mg
- Anhydrous Caffeine - 25 mg

Additives

Carmellose calcium, anhydrous silicic acid, gelatin, lactose, white soft sugar, starch, corn starch, aspartame (L-phenylalanine compound), vanillin and perfume

Storage and Handling Cautions

- (1) Store this drug in a cool, dry place away from sunlight and moisture.
- (2) Keep out of reach of children.
- (3) Do not transfer this drug to a different container (It may cause misuse or deterioration of quality).
- (4) When you use one sachet in several batches, seal the opened package by folding the edge during storage and use within 2 days.
- (5) Do not take this drug after the expiration date.

製造販売元  **エスエス製薬株式会社**
〒163-1488 東京都新宿区西新宿3-20-2
<https://www.ssp.co.jp/>

The marketing of this drug has been approved for domestic sales in accordance with the Japanese Pharmaceutical and Medical Devices Act (PMD Act.)