

Please read the instructions carefully before using.

エスタック®総合感冒

かぜの諸症状（のどの痛み・熱）の緩和に **第②類医薬品**

※ この医薬品は「使用上の注意」をよく読んでお使いください。

アレルギー体質の方は、必ず薬剤師、登録販売者にご相談ください。



S. TAC GENERAL COLD MEDICINE

Designated Class 2 OTC drug

General cold medicine

For cold of the whole family (5 years or older)

Film-coated tablets

- S. TAC GENERAL COLD MEDICINE is a general cold medicine for family use which the whole family with 5 years or older can use.
- It is a film-coated tablet easy to swallow with the formula which can work well for 11 symptoms of cold.

CAUTIONS FOR USE

CONTRAINDICATIONS

(Failure to observe the following may cause the present symptoms to worsen or produce adverse drug reactions and/or accidents.)

1. The following people must not take this drug:

- (1) Those with a previous allergic reaction to this drug or ingredients of the drug
- (2) Those who have ever suffered from asthma after taking this drug or other drugs for cold remedies or for relief of pain and fever

2. Do not take this drug in combination with any of the following drugs:

Other cold medicines, pain relievers/fever reducers, sedatives, cough and expectorant medicines, antihistamine-containing oral drugs (e.g. oral nasal inflammation drugs, motion sickness drugs, anti-allergy drugs, hypnotics and sedatives, etc.)

3. Do not drive or operate machinery after taking this drug.

(Sleepiness, etc. may occur.)

4. Do not drink alcohol before or after taking this drug.

5. Do not use this drug for a prolonged period of time.

WHEN to consult a doctor

1. The following people should consult a doctor, pharmacist or registered salesperson before taking this drug:

- (1) Those who are under the treatment of a doctor or dentist
- (2) Pregnant women or those who may become pregnant
- (3) Breastfeeding women
- (4) Elderly people

®: Trademark



SSP Co., Ltd.

- (5) Those with any previous allergic reactions to drugs or other things
- (6) Those with any of the following symptoms: High fever or urination difficulties
- (7) Those who were diagnosed with any of the following diseases:
Disrupted thyroid function, diabetes, heart disease, hypertension, liver disease, kidney disease, gastric and duodenal ulcers, glaucoma

2. The following symptoms may be adverse drug reactions. If any of these symptoms occur, STOP TAKING the drug immediately and consult a doctor, pharmacist or registered salesperson. Show them this leaflet.

Related areas	Symptoms
Skin	Rash, redness, itching
Digestive system	Nausea, vomiting, loss of appetite
Central nervous system	Dizziness/vertigo
Respiratory system	Breath shortness
Urinary system	Urination difficulties
Other	Excessive fall in temperature

In rare cases, the following serious symptoms could occur. In such an event, immediately consult a doctor.

Symptom's name	Symptoms
Shock (Anaphylaxis)	Itching of skin, hives, hoarseness, sneezing, itching of throat, tightness in the chest, palpitations, consciousness clouding, etc. which develop soon after taking the drug
Mucocutaneous ocular syndrome (Stevens-Johnson syndrome), Toxic Epidermal necrolysis, Acute generalized exanthematous pustulosis (AGEP)	Persistent or acute worsening of high fever, bloodshot eyes, eye discharge, lip erosion, sore throat, widespread rash, redness, redness of skin with small pimples, general malaise, loss of appetite, etc.
Drug-induced hypersensitivity syndrome	Redness of the skin over a wide range, generalized rash, fever, dullness, swelling of lymph nodes (neck, underarm, base of hip, etc.), etc.
Liver dysfunction	Fever, itching, rash, jaundice (skin and whites of the eyes turn yellow), brown urine, general malaise, loss of appetite, etc.
Kidney disorder	Fever, rash, reduction of urine output, generalized swelling, general malaise, joint pain (painful joints), diarrhea, etc.
Interstitial pneumonia	Shortness of breath, difficulty breathing when going up stairs or performing strenuous activity, dry cough, fever or other symptoms, which happen suddenly or continuously
Asthma	Wheezing or crowing sound when breathing breathlessness, etc.
Aplastic anemia	Purpura, nose bleeding, gum bleeding, fever, pale skin, pale mucosa, tiredness, palpitations, breathlessness, giddy feeling accompanied by queasy feeling, bloody urine, etc.
Agranulocytosis	Sudden high fever, chill, sore throat, etc.

3. You may develop the following symptoms after taking this drug. If the symptoms continue or worsen, stop taking the drug and consult a doctor, pharmacist or registered salesperson. Show them this leaflet.

Dry mouth or sleepiness

4. If the symptoms do not improve after 5 or 6 doses, stop taking this drug and consult a doctor, pharmacist or registered salesperson. Show them this leaflet.

Indications

Alleviation of various common cold symptoms (sore throat, runny nose, nasal congestion, fever, chills, headache, sneezing, cough, sputum, joint pain and muscle pain)

Dosage and Directions

Take the following dose 3 times a day, preferably within 30 minutes of a meal. This drug should be taken with either cold or lukewarm water.

Age	Adult (15 years or over)	11-14 years	5-10 years	Under 5 years
One dose	3 tablets	2 tablets	1 tablet	Do not take

<Cautions regarding dosage and directions>

- (1) Strictly observe the dosage and directions.
- (2) Children must be supervised by a parent or guardian when taking this drug.

Ingredients

Nine tablets (daily adult usage) contain:

- Acetaminophen - 900 mg
- Chlorpheniramine Maleate - 7.5 mg
- Dextromethorphan Hydrobromide Hydrate - 48 mg
- dl-Methylephedrine Hydrochloride - 60 mg
- Hesperidin - 45 mg
- Glycyrrhiza Extract - 187.5 mg (Equivalent to Glycyrrhiza - 750 mg)
- Powdered Ginger - 150 mg
- Anhydrous Caffeine - 75 mg

Additives

Croscarmellose sodium, anhydrous silicic acid, cellulose, lactose, hypromellose, povidone, macrogol, magnesium stearate, talc, titanium oxide, corn starch and red ferric oxide

Storage and Handling Cautions

- (1) Store this drug in a cool, dry place away from sunlight and moisture.
- (2) Keep out of reach of children.
- (3) Do not transfer this drug to a different container (It may cause misuse or deterioration of quality).
- (4) Tighten the bottle cap securely. If you do not replace the cap tightly, the tablet may become damaged by moisture. Additionally, do not touch tablets with wet hands.
- (5) The padding in the bottle is to prevent damage of the tablets during transportation. Discard it after opening as it is no longer necessary.
- (6) Do not take this drug after the expiration date.

製造販売元  **エスエス製薬株式会社**
〒163-1488 東京都新宿区西新宿3-20-2
<https://www.ssp.co.jp/>

The marketing of this drug has been approved for domestic sales in accordance with the Japanese Pharmaceutical and Medical Devices Act (PMD Act.)