

Please read the instructions carefully before using.

エスタック®総合 IB

第②類医薬品

※ この医薬品は「使用上の注意」をよく読んでお使いください。

アレルギー体質の方は、必ず薬剤師、登録販売者にご相談ください。



S.TAC GENERAL IB

Designated Class 2 OTC drug

Cold medicine

<Product Features>

- S.TAC GENERAL IB is a general cold medicine that works well from the beginning of catching a cold.
- It contains ibuprofen and herbal medicines (cinnamon bark and ginger). It is effective for the symptoms such as “chills, sore throat” etc. at the onset of a cold.
- It is also recommended as a household medicine.

CAUTIONS FOR USE

CONTRAINDICATIONS

(Failure to observe the following may cause the present symptoms to worsen or produce adverse drug reactions and/or accidents.)

1. The following people must not take this drug:

- (1) Those with a previous allergic reaction to this drug or ingredients of the drug
- (2) Those who have ever suffered from asthma after taking this drug or other drugs for cold remedies or for relief of pain and fever
- (3) Children under 15 years of age
- (4) Pregnant women whose due date is within 12 weeks

2. Do not take this drug in combination with any of the following drugs:

Other cold medicines, pain relievers/fever reducers, sedatives, cough and expectorant medicines, antihistamine-containing oral drugs (e.g. oral nasal inflammation drugs, motion sickness drugs, anti-allergy drugs, hypnotics and sedatives, etc.)

3. Do not drive or operate machinery after taking this drug.

(Sleepiness may occur.)

4. Breastfeeding women should not take this drug. If unavoidable, refrain from breastfeeding while taking this drug.

5. Do not drink alcohol before or after taking this drug.

6. Do not exceed 5 days of usage.

®: Trademark



SSP Co., Ltd.

WHEN to consult a doctor

1. The following people should consult a doctor, pharmacist or registered salesperson before taking this drug:

- (1) Those who are under the treatment of a doctor or dentist
- (2) Pregnant women or those who may become pregnant
- (3) Elderly people
- (4) Those with any previous allergic reactions to drugs or other things
- (5) Those with any of the following symptoms: High fever or urination difficulties
- (6) Those who were diagnosed with any of the following diseases:
Disrupted thyroid function, diabetes, heart disease, hypertension, liver disease, kidney disease, glaucoma, systemic lupus erythematosus, mixed connective-tissue disease, respiratory functional disorder, obstructive sleep apnoea syndrome (OSAS) or obesity
- (7) Those who have suffered from any of the following diseases:
Gastric and duodenal ulcers, ulcerative colitis or Crohn's disease

2. The following symptoms may be adverse drug reactions. If any of these symptoms occur, STOP TAKING the drug immediately and consult a doctor, pharmacist or registered salesperson. Show them this leaflet.

Related areas	Symptoms
Skin	Rash, redness, itching or purpura
Digestive system	Nausea, vomiting, loss of appetite, stomach discomfort, stomachache, oral ulcer, heartburn, heavy stomach feeling, gastrointestinal bleeding, abdominal pains, diarrhea or bloody stool
Central nervous system	Dizziness/vertigo
Circulatory system	Palpitations
Respiratory system	Breath shortness
Urinary system	Urination difficulties
Other	Blurred vision, ringing in the ears, swelling, nose bleeding, gum bleeding, uncontrolled bleeding, bleeding, back pain, excessive fall in temperature or malaise

In rare cases, the following serious symptoms could occur. In such an event, immediately consult a doctor.

Symptom's name	Symptoms
Shock (Anaphylaxis)	Itching of skin, hives, hoarseness, sneezing, itching of throat, tightness in the chest, palpitations, consciousness clouding, etc. which develop soon after taking the drug
Mucocutaneous ocular syndrome (Stevens-Johnson syndrome), Toxic Epidermal necrolysis	Persistent or acute worsening of high fever, bloodshot eyes, eye discharge, lip erosion, sore throat, widespread rash, redness, etc.

(Continued from the previous page)

Symptom's name	Symptoms
Liver dysfunction	Fever, itching, rash, jaundice (skin and whites of the eyes turn yellow), brown urine, general malaise, loss of appetite, etc.
Kidney disorder	Fever, rash, reduction of urine output, generalized swelling, general malaise, joint pain, diarrhea, etc.
Aseptic meningitis	Severe headache with neck muscle rigidity, fever, nausea, vomiting, etc. (These symptoms are mainly reported in patients undergoing treatment for systemic lupus erythematosus or mixed connective-tissue disease.)
Interstitial pneumonia	Shortness of breath, difficulty breathing when going up stairs or performing strenuous activity, dry cough, fever or other symptoms, which happens suddenly or continuously
Asthma	Wheezing or crowing sound when breathing breathlessness, etc.
Aplastic anemia	Purpura, nose bleeding, gum bleeding, fever, pale skin, pale mucosa, tiredness, palpitations, breathlessness, giddy feeling accompanied by queasy feeling, bloody urine, etc.
Agranulocytosis	Sudden high fever, chill, sore throat, etc.
Respiratory depression	Shortness of breath or difficulty breathing, etc.

3. You may develop the following symptoms after taking this drug. If the symptoms continue or worsen, stop taking the drug and consult a doctor, pharmacist or registered salesperson. Show them this leaflet.

Constipation, dry mouth or sleepiness

4. If the symptoms do not improve after 5 or 6 doses, stop taking the drug and consult a doctor, pharmacist or registered salesperson. Show them this leaflet.

Indications

Alleviation of various common cold symptoms (fever, chills, sore throat, cough, sputum, runny nose, nasal congestion, sneezing, headache, joint pain and muscle pain)

Dosage and Directions

Take the following dose 3 times a day, preferably within 30 minutes of a meal. This drug should be taken with either cold or lukewarm water.

Age	Adult (15 years or over)	Under 15 years
One dose	2 tablets	Do not take

<Cautions regarding dosage and directions>

Strictly observe the dosage and directions.



SSP Co., Ltd.

Ingredients

Six tablets contain (daily dose):

- Ibuprofen - 450 mg
- Powdered Ginger - 200 mg
- Powdered Cinnamon Bark - 400 mg
- Dihydrocodeine phosphate - 24 mg
- dl-Methylephedrine hydrochloride - 60 mg
- Chlorpheniramine maleate - 7.5 mg
- Anhydrous caffeine - 75 mg

Additives

Anhydrous silicic acid, cellulose, lactose, hydroxypropyl cellulose, hypromellose, macrogol, magnesium stearate, talc, titanium oxide

Storage and Handling Cautions

- (1) Store this drug in a cool, dry place away from sunlight and moisture.
- (2) Keep out of reach of children.
- (3) Do not transfer this drug to a different container as it may cause misuse or deterioration.
- (4) Tighten the bottle cap securely. If you do not replace the cap tightly, the tablet may become damaged by moisture. Additionally, do not touch tablets with wet hands.
- (5) The stuffing is to protect tablets against damage during shipping. Please discard it after opening.
- (6) Do not take this drug after the expiration date.

製造販売元



エスエス製薬株式会社
〒163-1488 東京都新宿区西新宿3-20-2
<https://www.ssp.co.jp/>

The marketing of this drug has been approved for domestic sales in accordance with the Japanese Pharmaceutical and Medical Devices Act (PMD Act.)