

Please read the instructions carefully before using.

# アネロン「ニスキャップ」

乗物酔い薬 第②類医薬品

※ この医薬品は、薬剤師、登録販売者に相談の上、  
「使用上の注意」をよく読んでお使いください。



ANERON"NYSCAPS"

Designated Class 2 OTC drug

## Motion sickness drug

- ANERON"NYSCAPS" is a capsule medicine that is effective for prevention and alleviation of nausea, dizziness and headache caused by motion sickness.
- It is a long-acting medicine that works by once daily dosing 30 minutes before departure.
- It alleviates nausea, dizziness and headache caused by motion sickness when you take soon after feeling of motion sickness.

## CAUTIONS FOR USE

### CONTRAINDICATIONS

(Failure to observe the following may cause the present symptoms to worsen or produce adverse drug reactions and/or accidents.)

- 1. The following people must not take this drug:**  
Children under 15 years of age.
- 2. Do not take this drug in combination with any of the following drugs:**  
Other motion sickness drugs, cold remedies, pain relievers/fever reducers, sedatives, cough and expectorant medicines, gastrointestinal pain reliever and antispasmodics, antihistamine-containing oral drugs (e.g. oral nasal inflammation drugs, anti-allergy drugs, etc.)
- 3. Do not drive or operate machinery after taking this drug.**  
(Sleepiness, blurred vision or excessive sensitivity to light may occur.)

### WHEN to consult a doctor

- 1. The following people should consult a doctor, pharmacist or registered salesperson before taking this drug:**
  - (1) Those who are under the treatment of a doctor
  - (2) Pregnant women or those who may become pregnant
  - (3) Elderly people
  - (4) Those with any previous allergic reactions to drugs or other things
  - (5) Those with any of the following symptoms: Urination difficulties
  - (6) Those who were diagnosed with any of the following diseases:  
Glaucoma or heart disease

- 2. The following symptoms may be adverse drug reactions. If any of these symptoms occur, STOP TAKING the drug immediately and consult a doctor, pharmacist or registered salesperson. Show them this leaflet.**

Related areas	Symptoms
Skin	Rash, redness, itching
Central nervous system	Headache
Circulatory system	Palpitations
Urinary system	Urination difficulties
Other	Hot flushes on face or excessive sensitivity to light

- 3. You may develop the following symptoms after taking this drug. If the symptoms continue or worsen, stop taking the drug and consult a doctor, pharmacist or registered salesperson. Show them this leaflet.**

Dry mouth, constipation, diarrhea, sleepiness or blurred vision.

## Indications

Prevention and alleviation of nausea, dizziness and headache caused by motion sickness.

## Dosage and Directions

Take the following dose once a day. For the prevention of motion sickness, take this drug 30 minutes before departure. This drug should be taken with water either cold or lukewarm.

Age	Adult (15 years or over)	Under 15 years
One dose	One capsule	Do not take

### <Cautions regarding dosage and directions>

- (1) Strictly observe the dosage and directions.
- (2) You can take this drug regardless before or after a meal.

## Ingredients

One capsule contains:

- Pheniramine Maleate - 30 mg
- Ethyl Aminobenzoate - 50 mg
- Scopolamine Hydrobromide Hydrate - 0.2 mg
- Anhydrous Caffeine - 20 mg
- Pyridoxine Hydrochloride (Vitamin B<sub>6</sub>) - 5 mg

## Additives

Silicon dioxide, gelatin, cellulose, white soft sugar, hydroxypropyl cellulose, ethylcellulose, glycerol esters of fatty acids, talc, corn starch, methacrylic acid copolymer L, sodium lauryl sulfate, propyl gallate, Vitamin B<sub>2</sub>, red No.3, yellow No.5 and blue No.1.

## Advice for People Who Tend to Have Motion Sickness

---

- Sleep well at night before departure
- Keep a good digestive health by eating foods easily digestible
- Stay at a seat that moves less in relaxed manner
- Distract your feeling by viewing the scenery or chatting, playing games
- The drug is effective when you take before departure

## Storage and Handling Cautions

---

- (1) Store this drug in a cool, dry place away from sunlight and moisture.
- (2) Keep out of reach of children.
- (3) Do not transfer this drug to a different container (It may cause misuse or deterioration of quality).
- (4) Do not take this drug after the expiration date.

製造販売元  **エスエス製薬株式会社**  
〒163-1488 東京都新宿区西新宿3-20-2  
<https://www.ssp.co.jp/>

The marketing of this drug has been approved for domestic sales in accordance with the Japanese Pharmaceutical and Medical Devices Act (PMD Act.)